



GLASS

The detection and reporting of colistin resistance

Second edition

Global Antimicrobial Resistance and
Use Surveillance System (GLASS)

GLASS

The detection and reporting of colistin resistance

Second edition

Global Antimicrobial Resistance and
Use Surveillance System (GLASS)



GLASS: the detection and reporting of colistin resistance, second edition

ISBN 978-92-4-001904-1 (electronic version)

ISBN 978-92-4-001905-8 (print version)

© World Health Organization 2021

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. GLASS: the detection and reporting of colistin resistance, second edition. Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IGO.

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <http://www.who.int/about/licensing>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Design and layout by 400 Communications Limited.

Contents

Acknowledgements	iv
Acronyms and abbreviations	v
Summary	vi
1 Introduction	1
<hr/>	
2 Acquired resistance to colistin	3
2.1 Mutational colistin resistance	3
2.2 Transferrable colistin resistance	3
<hr/>	
3 Laboratory detection of colistin resistance	5
3.1 Phenotypic methods	5
3.2 Genotypic methods	7
<hr/>	
4 Surveillance strategies	8
<hr/>	
5 References	10
<hr/>	

Acknowledgements

WHO thanks the following authors and contributors at WHO collaborating centres:

Valeria Bortolaia; Alejandra Corso; Andrey Dekhnich; Rene S. Hendriksen; Roman Kozlov; Monica Lahra; Jean Patel; Wantana Paveenkittiporn; Olga Perovic; Ana Rita Rebelo and Neil Woodford.

Contributions by staff at the WHO regional offices

Sheick Oumar Coulibaly, Walter Fuller, Laetitia Gahimbare for the Regional Office for Africa (AFRO); Marcelo Galas and Ramon Pardo Pilar at the Regional Office for the Americas (PAHO); Mona Elshokry and Franciscus Konings at the Regional Office for the Eastern Mediterranean (EMRO); Danilo Lo Fo Wong at the Regional Office for Europe (EURO); Aparna Singh Shah and Sirenda Vong at the Regional Office for South East Asia (SEARO); and Socorro Escalante, Raynal Squires and Babatunde Olowokure at the Regional Office for the Western Pacific (WPRO).

External reviewers

Anette M. Hammerum (Statens Serum Institute, Denmark); Rumina Hasan (Aga Khan University, Pakistan); Susan Van Den Hof and Ed Kuijper (National Institute for Public Health and the Environment, RIVM, Netherlands); Gunnar Kahlmeter and Erika Matuschek (European Committee on Antimicrobial Susceptibility Testing, EUCAST); Maria Karlsson and Joseph Lutgring (Centers for Disease Control and Prevention, USA); and Gregory Tyson (Food and Drug Administration, USA).

WHO staff

Jorge Raul Matheu Alvarez; Sebastien Cognat; Sergey Romualdovich Eremin; Sapna Manglani; Christopher Oxenford and Carmem Lucia Pessoa-Silva.

Developer group

Jorge Raul Matheu Alvarez; Rene S. Hendriksen; Sapna Manglani.

Executive group

Rene S. Hendriksen; Carmem Lucia Pessoa-Silva.

Editing

Elisabeth Heseltine.

Financial support

The Government of the United States of America.

Acronyms and abbreviations

AMR	Antimicrobial resistance
CAT	Colistin Agar Test
CBDE	Colistin Broth Disk Elution
CLSI	Clinical and Laboratory Standards Institute
EUCAST	European Committee on Antimicrobial Susceptibility Testing
GLASS	Global Antimicrobial Resistance Surveillance System
MIC	Minimum inhibitory concentration
PCR	Polymerase chain reaction

Summary

The scope of this technical note is to review current methods for the detection of colistin resistance and to provide a framework for its investigation. The document highlights the critical distinction between phenotypic detection of colistin resistance and genotypic detection of specific colistin resistance mechanisms, such as *mcr* and chromosomal mutations. Colistin resistance in Enterobacterales and *Acinetobacter baumannii* is included in the WHO Global Antimicrobial Resistance Surveillance System (GLASS). Currently, however, reliable tests for phenotypic detection of colistin resistance for clinical and surveillance purposes are not widely available.

The technical note describes existing phenotypic methods for detecting colistin resistance, genotypic methods for detecting specific colistin resistance mechanisms and surveillance strategies for monitoring colistin resistance.

This is a rapidly evolving field with new resistance genes being identified regularly and new methods for phenotypic resistance testing being described, hence this document will be updated, as needed, to reflect these developments.

This is the second edition of *The detection and reporting of colistin resistance* that was first published in December 2018 (<https://apps.who.int/iris/bitstream/handle/10665/277175/WHO-WSI-AMR-2018.4-eng.pdf>). This new edition includes information on additional *mcr* genes which have since been identified. It also includes updated information in the section on the laboratory related detection of colistin resistance. Additional technical guidance is provided on phenotypic methods, including information on some recently published novel methods. The section on genotypic methods includes updated technical guidance and tools.

Key Points

- In humans, colistin is generally used as a last resort to treat infections with highly resistant Gram-negative bacteria.
- Resistance to colistin is ongoing, with new resistance genes emerging.
- Acquired colistin resistance mechanisms are both plasmid-mediated and chromosomal.
- Phenotypic testing for colistin susceptibility has inherent technical difficulties, and only broth microdilution is currently recommended.
- Genotypic testing is limited to the detection of acquired resistance genes, so a negative result in a PCR test cannot be used to predict susceptibility to colistin.
- Surveillance of colistin resistance in *Enterobacterales* and *Acinetobacter baumannii* is included in the WHO GLASS.
- This is a rapidly evolving field, and this document will be updated as needed.

01 Introduction

Colistin belongs to a group of antimicrobial agents known as polymyxins, which were originally isolated from the spore-forming soil organism *Paenibacillus polymyxa*. Molecules in this group are polymyxins A, B, C, D and E, of which only polymyxin E (colistin) and polymyxin B are used clinically in humans.

Colistin is a mixture of polymyxin E1 and E2, two bactericidal pentacationic lipopeptides whereas polymyxin B is composed of polymyxins B1, B1-I, B2, B3, and B6. Polymyxin B has bactericidal action against almost all Gram-negative bacteria.

The mode of action of colistin is not fully elucidated but involves binding to lipopolysaccharides and phospholipids in the outer membrane of Gram-negative organisms, which results in membrane disruption and cell death. Colistin is active against a wide variety of Gram-negative bacteria and is not active against Gram-positive bacteria, which lack an outer membrane.

Colistin is used in both human and veterinary medicine. In humans, colistin is generally used to treat infections with multidrug-resistant, extensively drug-resistant and pan drug-resistant bacteria (1). It is usually administered by injection or inhalation (the latter, for example, for patients with cystic fibrosis) as the sodium salt of colistin methanesulfonate, which is an inactive prodrug (2,3). It is considered less toxic than colistin sulfate, which is used orally (with very limited absorption) or topically as for selective gut decontamination in patients with hematological diseases or at Intensive Care Units (4).

In veterinary medicine, colistin has been widely used in various food-producing animals (broiling and laying hens, pigs, calves, beef cattle, dairy cattle, meat- and milk-producing sheep and goats, rabbits and fish in aquaculture) for the prevention and treatment of infections caused by Enterobacteriaceae and other Gram-negative bacteria and for growth promotion. In veterinary medicine, colistin is usually administered orally (5). The wide use of colistin in food-producing animals is considered to have contributed extensively to the emergence of resistance to this drug (5).

With the spread of highly resistant bacteria such as carbapenemase-producing Enterobacterales and multi-drug resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii*, colistin has been used increasingly as an agent of last resort. The emergence of colistin resistance in multidrug-resistant bacteria is therefore a significant clinical and public health concern (6-9).

The spectrum of colistin (Table 1) includes activity in vitro against *Acinetobacter* spp., *Pseudomonas aeruginosa*, *Klebsiella* spp., *Enterobacter* spp., *Escherichia coli*, *Salmonella* spp. (MIC values are often higher than other species), *Shigella* spp., *Citrobacter* spp., *Yersinia pseudotuberculosis* and *Haemophilus influenzae*. Furthermore, colistin has considerable activity against *Stenotrophomonas* spp., *Vibrionaceae* and *Bordetella* spp., whereas it is not active against some Gram-negative aerobic bacilli, including *Burkholderia mallei*, *Burkholderia cepacia*, *Proteus* spp., *Providencia* spp., *Morganella morganii*, *Serratia* spp., *Edwardsiella* spp. and *Brucella* spp. (10-13), Gram-negative and Gram-positive aerobic cocci, Gram-positive aerobic bacilli, some anaerobes, fungi and parasites (4,14).

Table 1. Activity of colistin against selected pathogenic bacteria

ORGANISM	NATURAL STATE (RESISTANT OR SUSCEPTIBLE TO POLYMYXINS)
<i>Escherichia coli</i>	Susceptible
<i>Klebsiella</i> spp.	Susceptible
<i>Enterobacter</i> spp.	Susceptible
<i>Pseudomonas aeruginosa</i>	Susceptible
<i>Acinetobacter</i> spp.	Susceptible
<i>Salmonella</i> spp.	Susceptible (MIC values are often higher than in other species)
<i>Shigella</i> spp.	Susceptible
<i>Pasteurella</i> spp.	Susceptible
<i>Haemophilus</i> spp.	Susceptible
<i>Aeromonas</i> spp.	Susceptible
<i>Proteus</i> spp.	Intrinsically resistant
<i>Providencia</i> spp.	Intrinsically resistant
<i>Morganella morganii</i>	Intrinsically resistant
<i>Serratia</i> spp.	Intrinsically resistant
<i>Brucella</i> spp.	Intrinsically resistant
<i>Neisseria</i> spp.	Intrinsically resistant
<i>Chromobacterium</i> spp.	Intrinsically resistant
<i>Burkholderia</i> spp.	Intrinsically resistant
Gram-positive bacteria	All Gram-positive bacteria are intrinsically resistant to polymyxins because of the absence of an outer membrane.

02

Acquired resistance to colistin

2.1 Mutational colistin resistance

Acquired resistance to polymyxins in naturally susceptible species is usually the result of modifications of the bacterial cell surface, including alterations to the lipopolysaccharide structure. Other reported resistance mechanisms include shedding of capsular polysaccharides, which trap or bind polymyxins, as found in some isolates of *Klebsiella pneumoniae* (15). In *A. baumannii*, two primary mechanisms of chromosomally mediated colistin resistance have been described. The first is due to loss of lipopolysaccharide production and the second to modification of the system that allows bacteria to respond to environmental conditions, affecting lipid modification and reducing bacterial membrane permeability (15,16). Resistance arising from chromosomal mutations is not usually transferred horizontally among bacteria (17).

2.2 Transferable colistin resistance

A plasmid-mediated gene that confers resistance to colistin, *mcr-1*, was first reported in late 2015 in *E. coli* isolates from food animals and their meat collected in China during 2011–2014 and in *E. coli* and *K. pneumoniae* isolates collected from human patients in China in 2014 (18). Although colistin had been used extensively in food-producing animals globally for many years, it was not available for human clinical use in China until 2017, suggesting that the selection pressure for the spread of *mcr-1* was driven by veterinary use of colistin (19).

Shortly after this first description, it was observed that the *mcr-1* gene could be found globally in *K. pneumoniae*, *E. coli*, *Enterobacter cloacae* and *Salmonella* spp. isolates of animal, environmental and human origin (18, 20–26). The global spread of *mcr-1* predated its recognition as a colistin resistance gene and was probably also facilitated by human travel, as suggested by the finding of the gene in enteric bacteria from travellers returning to Europe after visiting countries with a high prevalence of *mcr-1* in South America, Asia and Africa (19, 27).

After identification of *mcr-1*, wide scientific attention led to recognition of multiple *mcr-1* variants (28-37) and ten additional (at the time of writing) *mcr* gene families (38). The latter include *mcr-2* in *E. coli* and *Salmonella* spp. from pigs, cattle and chickens in Belgium (39, 40), *mcr-3* in human and animal *E. coli*, *Proteus* spp. and *Aeromonas* spp. in countries in Asia, Europe and South America (41), *mcr-4* in *E. coli* and *Salmonella enterica* serovar Typhimurium from pigs and humans in Belgium, Italy and Spain (42, 43), *mcr-5* in poultry and poultry meat isolates of *S. enterica* serovar Paratyphi B d-tartrate fermenting and in porcine *E. coli* in Germany (44, 45), *mcr-6* in *Moraxella* sp. isolates from pigs in the United Kingdom (36), *mcr-7* in *K. pneumoniae* from chickens in China (46, 47), *mcr-8* in *K. pneumoniae* from humans and pigs in China (48), *mcr-9* in a *Salmonella Typhimurium* strain isolated from a human patient in Washington State, USA (49) and *mcr-10* in a clinical strain of *Enterobacter roggkampii* isolated in China (50). It is noteworthy to mention that the *S. Typhimurium* strain in which *mcr-9* originally was detected was phenotypically susceptible to colistin according to EUCAST with an MIC of 2 Mg/L. The co-occurrence of more than one of these genes has been reported in *E. coli* and *Salmonella* spp. isolates of human and animal origin in Europe and China, such as the simultaneous presence of *mcr-1* and *mcr-3* and of *mcr-1* and *mcr-4* (51-55).

Bacteria isolated from food-producing animals to date appear to carry the *mcr* genes more frequently than bacteria isolated from humans, which is probably a consequence of the selective pressure exerted by the wide use of colistin in veterinary practice (15). Importantly, *mcr*-mediated colistin resistance can be transferred among bacterial strains, species and genera. To limit further dissemination of such genes, accurate identification of colistin-resistant, *mcr*-encoding isolates is of critical importance.

03 Laboratory detection of colistin resistance

3.1 Phenotypic methods

Testing for antimicrobial susceptibility by phenotypic methods is currently the cornerstone of AMR surveillance and is necessary to determine the occurrence of colistin resistance within GLASS. Phenotypic testing for antimicrobial susceptibility to colistin is, however, complex. The technical difficulties in phenotypic testing for colistin susceptibility include poor diffusion of polymyxins through agar, which compromises the performance of both disc diffusion and gradient diffusion methods, and the tendency of polymyxins to bind to the surface of plastics (e.g. the plastic of microtitre trays used for broth microdilution and of pipette tips) (56).

International standard-setting organizations such as the Clinical and Laboratory Standards Institute (CLSI)¹ and the European Committee on Antimicrobial Susceptibility Testing (EUCAST)² formed a joint polymyxin breakpoints working group in March 2016, which recommended only broth microdilution for testing of colistin susceptibility. Thus, the susceptibility of Enterobacteriaceae, *P. aeruginosa* and *Acinetobacter* spp. to colistin should be tested according to the International Organization for Standardization standard broth microdilution method (ISO 20776-1), in which cation-adjusted Mueller-Hinton broth is used. No additives may be included in any part of testing (in particular no polysorbate-80 or other surfactants), because, for example, polysorbate 80 can act synergistically with polymyxins and artificially lower the minimum inhibitory concentration (MIC) (57). Trays should be made of plain polystyrene, and sulfate salts of polymyxins must be used (the methane sulfonate derivative of colistin cannot be used as it is an inactive pro-drug that breaks down slowly in solution). A review of the performance of different broth microdilution-based systems used for colistin susceptibility testing is available (58). CLSI has recently included two additional methods, Colistin Broth Disk Elution (CBDE) and Colistin Agar Test (CAT) as acceptable for detection of colistin resistance (59). Broth microdilution remains the only approved and recommended method for polymyxin B.

Quality control is essential in phenotypic detection of colistin resistance. To ensure that the method used provides reliable results, inclusion of a strain with low-level colistin resistance (such as *mcr-1*-positive *E. coli* NCTC 13846) is recommended by EUCAST (60).

Phenotypic results can be interpreted and reported as either the epidemiological cut-offs (according to EUCAST or CLSI), which define wild-type and non-wild-type populations, or as clinical breakpoints, which define clinical susceptibility and resistance (Table 2) (61-63).

¹ <https://clsi.org/>

² <http://www.eucast.org/>

Table 2. EUCAST and CLSI clinical break points for colistin

BACTERIA	SUSCEPTIBLE	INTERMEDIATE	RESISTANT
Enterobacteriaceae			
EUCAST	=< 2	--	>= 4
CLSI	-	=< 2	>= 4
<i>Acinetobacter</i> spp.*			
EUCAST	=< 2	--	>= 4
CLSI	-	=< 2	>= 4
<i>Pseudomonas aeruginosa</i>			
EUCAST	=< 2	--	>= 4
CLSI	-	=< 2	>= 4

* CLSI breakpoints apply only for *A. baumannii* complex.

Clinical trials, including ones in which new antibiotics are compared, are providing new data on the clinical effectiveness of colistin for the treatment of infections caused by Gram-negative bacteria. According to the clinical outcomes, colistin appears to be a poor mono-therapeutic option, especially when other drugs are available (64, 65). In particular, the dosing of colistin is complex, as its pharmacokinetics differs widely among patients (66). Therefore, further review of clinical outcomes is necessary to determine whether currently established breakpoints and epidemiological cut-off values accurately inform clinical decisions. See CLSI and EUCAST documents for the latest information.

Testing for antimicrobial susceptibility by broth microdilution requires greater laboratory capacity than standard disc diffusion and gradient diffusion methods and other phenotypic methods. Nevertheless, other methods are either insufficiently accurate or have not yet undergone rigorous multicentre evaluation with large collections of bacteria, including different species (i.e. most Enterobacteriaceae and Gram-negative non-fermenters), that express all known colistin resistance mechanisms. Thus, these methods cannot yet be recommended as best practice.

The methods that appear promising but are still to undergo the rigorous evaluation described above are: (i) agar-based screening media for bacteria suspected to be colistin-resistant (67-69), which should always be followed by further investigation of the colonies growing on such media by broth microdilution and/or molecular methods; (ii) the Rapid Polymyxin NP test (70); (iii) Colispot (58); and (iv) disc prediffusion (71). The performance of these methods depends on the bacterial species and the colistin resistance mechanisms (72).

EUCAST considers MIC-determination by broth microdilution the only valid method for colistin susceptibility testing. Alternative methods already tested by EUCAST failed to meet reasonable standards (72). CLSI considers MIC determination by broth microdilution, and the new proposed broth disk elution and agar dilution methods, as acceptable for colistin testing in Enterobacterales and *Pseudomonas aeruginosa* (73, 74). In contrast, broth microdilution is the only CLSI approved method for colistin and polymyxin B testing in *Acinetobacter* spp. Colistin broth disk elution and agar dilution tests are considered provisional until additional data are evaluated. Alternative methods like colistin disk elution drop test and colistin spot test have recently been proposed (75). These methods were validated with a limited number of brands of disks and/or media.

Laboratories need to take responsibility for validating the use of alternative methods. The recognized need for simpler, more reliable in vitro testing of colistin susceptibility and for methods to selectively identify colistin-resistant, *mcr*-positive isolates is driving technical development, and improved phenotypic testing products are expected to become available.

3.2 Genotypic methods

During its early implementation stage, GLASS only collects data on antimicrobial susceptibility of target pathogens, acquired by phenotypic testing (76); it does not yet collect data on the distribution of specific resistance mechanisms. Nevertheless, genotypic surveillance can be used as a basis for selection of isolates for subsequent phenotypic testing for antimicrobial susceptibility and the results submitted to GLASS, or it can be used as a supplementary investigation of resistance mechanisms for isolates with unusual phenotypic resistance(s). It is noteworthy to mention a few limitations, that the presence of transposons in association with a *mcr*-gene might hamper expression of the gene in phenotypic testing (77) as well as potential biased estimates of susceptibility that might lead to an overestimate of phenotypic resistance. Such studies can help in understanding the epidemiology of AMR and the relative contributions of the human clinical, animal health and agricultural sectors to the problem of AMR. They may also improve reporting to the Emerging Antimicrobial Resistance module in GLASS (the GLASS-EAR system).

Genotypic results should be reported as the presence or absence of the specified colistin resistance mechanism or gene. At present, there is insufficient understanding of acquired colistin resistance mechanisms to design a molecular test that would be sufficiently sensitive or specific to be recommended as best practice. In particular, genotypic methods would probably not detect all the diverse chromosomal mutations known to be responsible for most phenotypic colistin resistance in clinical settings (15) but would be limited to the detection of acquired colistin resistance genes. A few published and/or commercially available colistin molecular tests are available for detecting transferrable colistin resistance, including two multiplex polymerase chain reactions (PCR) for screening for the presence of *mcr-1* to *mcr-5* (78) and *mcr-6* to *mcr-9*, respectively (79).

It should be noted that a negative result in a PCR molecular test cannot be used to predict susceptibility to colistin, because the test cannot exclude the presence of chromosomal mechanisms of resistance or even of novel *mcr* genes that are not included in the test. As evidence of this limitation, high rates of colistin resistance are reported among strains of *K. pneumoniae* producing carbapenemase in Brazil and Italy but which lack *mcr* genes (80). In these settings, a negative PCR result for *mcr* genes would have poor predictive value for a colistin-susceptible phenotype. Whole-genome sequencing would allow screening for *mcr* genes and known chromosomal mutations that confer colistin resistance (81). Bioinformatics analysis could be conducted with, among others (82), the Center for Genomic Epidemiology Web tools (83), and ResFinder 4 (81, 84). Although the sensitivity and negative predictive value would be affected by inclusion of strains with novel mechanisms of resistance, this is the most comprehensive method for detecting all currently known putative colistin resistance mechanisms. It would also enable retrospective analysis of sequencing data as new resistance mechanisms are described.

Whatever the molecular method used, it is critically important to ensure that the PCR either detects all currently known *mcr* genes or that the databases used to impute resistance mechanisms from whole-genome sequencing data are up to date.

As our understanding of colistin resistance mechanisms improves, so will the concordance between phenotypic and genotypic test results. As for many other classes of antimicrobial agents, molecular testing may eventually offer an alternative to phenotypic testing for the surveillance of colistin resistance. Nevertheless, if the results are intended to guide clinical management, inference of phenotype based solely on a genotypic result may be valid only when the genotypic result is positive (i.e. mechanisms or genes detected), with a cautionary note that the resistance phenotype is likely but not guaranteed. If the results of the genotypic test are negative, no inference should be drawn about phenotype.

04 Surveillance strategies

In view of the many technical challenges to colistin susceptibility testing, clear strategies are required for clinical diagnosis, sampling and testing for surveillance. In local health care settings, colistin resistance should be tested if the recommended technology is available, and samples should be sent to laboratories with the necessary technical capacity if the technology is not available. The cost of shipping isolates is an important aspect to be considered in confirmatory testing. Testing should be performed on the following, recognising that intrinsically colistin-resistant genera are excluded:

- carbapenem-resistant Enterobacteriaceae
- other Gram-negative bacteria resistant to carbapenems and,
- when treatment with colistin must be considered

Health care facilities with a high prevalence or outbreaks of infections caused by carbapenem-resistant and other multidrug-resistant Enterobacteriaceae and/or non-fermenting bacteria such as *Acinetobacter baumannii* and *P. aeruginosa*, could consider conducting periodic surveys to assess the occurrence of colistin resistance to inform local antimicrobial use policies.

At the national level, at least one AMR reference laboratory should be established in each member state to monitor colistin resistance reliably, with appropriate controls and standardized methods. Such an AMR reference laboratory could perform confirmatory testing upon request from local laboratories. Samples for confirmatory testing should be selected carefully, with priority given to isolates suspected to have acquired transferrable (*mcr*-mediated) colistin resistance. A reasonable strategy for detecting *mcr*-mediated colistin resistance would be to screen for phenotypic resistance (for example, using selective agar plates, CHROMID® Colistin R agar, Superpolymyxin or CHROMagar COL-APSE) (67, 68, 85, 86) followed by PCR to detect one or more of the *mcr* genes. Such molecular testing could be performed at a reference laboratory or at a WHO Collaborating Centre.

Ideally, colistin susceptibility testing should be included in national plans for AMR surveillance. Countries with strong national AMR surveillance programmes and laboratory capacity could extend them to include monitoring for colistin resistance in bacteria isolated from food-producing animals and from food and environmental samples, at least periodically.

Currently, the detection of colistin resistance is too technically complicated to recommend as a phenotypic screening method for resource-limited settings. It is therefore essential that laboratories store isolates suspected of being colistin-resistant and collaborate with AMR reference laboratories and/or WHO Collaborating Centres to confirm colistin resistance and determine the possible presence of *mcr* genes.

The GLASS protocol (73) for sample-based surveillance is recommended for colistin susceptibility testing. In sample-based surveillance, case-finding is focused on priority specimens sent routinely to laboratories for clinical purposes. In addition, epidemiological information should be clearly linked to microbiological information for each sample and the standard AMR indicators recorded for positive cultures. The sample-based approach allows acquisition of information on the proportion of susceptible, intermediate and resistant isolates and the total number of patients sampled, making it possible to deduce the rates or frequencies of specific types of resistance in the tested population. Information on the incidence of this type of resistance among tested patients could provide an indication of the extent of colistin resistance in the population. The sample population must be carefully defined and described in any report in order not to overestimate resistance rates if only highly resistant strains are tested.

05

References

1. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* 2012;18(3):268–81.
2. Vardakas KZ, Voulgaris GL, Samonis G, Falagas ME. Inhaled colistin monotherapy for respiratory tract infections in adults without cystic fibrosis: a systematic review and meta-analysis. *Int J Antimicrob Agents.* 2018;51(1):1–9.
3. Barry PJ, Jones AM. New and emerging treatments for cystic fibrosis. *Drugs.* 2015;75(11):1165–75.
4. Falagas ME, Kasiakou SK. Colistin: the revival of polymyxins for the management of multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis.* 2005;40(9):1333–41.
5. Catry B, Cavaleri M, Baptiste K, Grave K, Grein K, Holm A, et al. Use of colistin-containing products within the European Union and European Economic Area (EU/EEA): development of resistance in animals and possible impact on human and animal health. *J Antimicrob Agents.* 2015; 46(3):297–306.
6. Jeannot K, Bolard A, Plésiat P. Resistance to polymyxins in Gram-negative organisms. *Int J Antimicrob Agents.* 2017;49(5):526–35.
7. Srinivas P, Rivard K. Polymyxin resistance in Gram-negative pathogens. *Curr Infect Dis Rep.* 2017;19(11):38.
8. Tumbarello M, Trecarichi EM, De Rosa FG, Giannella M, Giacobbe DR, Bassetti M, et al. Infections caused by KPC-producing *Klebsiella pneumoniae*: differences in therapy and mortality in a multicentre study. *J Antimicrob Chemother.* 2015;70(7):2133–43.
9. Tsuji BT, Pogue JM, Zavascki AP, et al. International Consensus Guidelines for the Optimal Use of the Polymyxins: Endorsed by the American College of Clinical Pharmacy (ACCP), European Society of Clinical Microbiology and Infectious Diseases (ESCMID), Infectious Diseases Society of America (IDSA), International Society for Anti-infective Pharmacology (ISAP), Society of Critical Care Medicine (SCCM), and Society of Infectious Diseases Pharmacists (SIDP). *Pharmacotherapy.* 2019;39(1):10-39. doi:10.1002/phar.2209
10. Muyembe T, Vandepitte J, Desmyter J. Natural colistin resistance in *Edwardsiella tarda*. *Antimicrob Agents Chemother.* 1973; 4(5):521–4.
11. Loutet SA, Valvano MA. Extreme antimicrobial Peptide and polymyxin B resistance in the genus *Burkholderia*. *Front Microbiol.* 2011; 2:159.
12. Biswas S, Brunel JM, Dubus JC, Reynaud-Gaubert M, Rolain JM. Colistin: an update on the antibiotic of the 21st century. *Expert Rev Anti Infect Ther.* 2012 Aug; 10(8): 917-34. doi: 10.1586/eri.12.78.

13. Samonis G, Korbila IP, Maraki S, Michailidou I, Vardakas KZ, Kofteridis D, et al. Trends of isolation of intrinsically resistant to colistin *Enterobacteriaceae* and association with colistin use in a tertiary hospital. *Eur J Clin Microbiol Infect Dis*. 2014; 33(9):1505–10.
14. Landman D, Georgescu C, Martin DA, Quale J. Polymyxins revisited. *Clin Microbiol Rev*. 2008;21(3):449–65.
15. Olaitan AO, Morand S, Rolain JM. Mechanisms of polymyxin resistance: acquired and intrinsic resistance in bacteria. *Front Microbiol*. 2014; 5:643.
16. Ahmed SS, Alp E, Hopman J, Voss A. Global epidemiology on colistin resistant *Acinetobacter baumannii*. *J Infect Dis Ther*. 2016;4(4):287.
17. McPhee JB, Lewenza S, Hancock RE. Cationic antimicrobial peptides activate a two-component regulatory system, PmrA-PmrB, that regulates resistance to polymyxin B and cationic antimicrobial peptides in *Pseudomonas aeruginosa*. *Mol Microbiol*. 2003; 50(1):205–17.
18. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism *MCR-1* in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016; 16:161–8.
19. Skov RL, Monnet DL. Plasmid-mediated colistin resistance (*mcr-1* gene): three months later, the story unfolds. *EuroSurveill*. 2016;21(9):30155.
20. Al-Tawfiq JA, Laxminarayan R, Mendelson M. How should we respond to the emergence of plasmid-mediated colistin-resistance in humans and animals. *Int J Infect Dis*. 2017; 54:77–84.
21. Coetzee J, Corcoran C, Prentice E, Moodley M, Mendelson M, Poirel L, et al. Emergence of plasmid-mediated colistin resistance (*MCR-1*) among *Escherichia coli* isolated from South African patients. *S Afr Med J*. 2016;106(5):35–6.
22. Gu DX, Huang YL, Ma JH, Zhou HW, Fang Y, Cai JC, et al. Detection of colistin resistance gene *mcr-1* in hypervirulent *Klebsiella pneumoniae* and *Escherichia coli* isolates from an Infant with diarrhea in China. *Antimicrob Agents Chemother*. 2016;60(8):5099–100.
23. Quan J, Li X, Chen Y, Jiang Y, Zhou Z, Zhang H, et al. Prevalence of *mcr-1* in *Escherichia coli* and *Klebsiella pneumoniae* recovered from bloodstream infections in China: a multicentre longitudinal study. *Lancet Infect Dis*. 2017;17(4):400–10.
24. Shen Z, Wang Y, Shen Y, Shen J, Wu C. Early emergence of *mcr-1* in *Escherichia coli* from food-producing animals. *Lancet Infect Dis*. 2016;16(3):293.
25. Wong SCY, Tse H, Chen JHK, Cheng VCC, Ho PL, Yeun KY. Colistin-resistant *Enterobacteriaceae* carrying the *mcr-1* gene among patients in Hong Kong. *Emerg Infect Dis*. 2016;22(9):1667–9.
26. Doumith M, Godbole G, Ashton P, Larkin L, Dallman T, Day M, et al. Detection of the plasmid-mediated *mcr-1* gene conferring colistin resistance in human and food isolates of *Salmonella enterica* and *Escherichia coli* in England and Wales. *J Antimicrob Chemother*. 2016;71(8):2300–5.
27. Arcilla MS, Van Hattem JM, Matamoros S, Melles DC, Penders J, De Jong MD, et al. Dissemination of the *mcr-1* colistin resistance gene. *Lancet Infect Dis*. 2016;16(2):147–9.
28. Di Pilato V, Arena F, Tascini C, Cannatelli A, Henrici De Angelis L, Fortunato S, et al. *mcr-1.2*, a new *mcr* variant carried on a transferable plasmid from a colistin-resistant KPC carbapenemase-producing *Klebsiella pneumoniae* strain of sequence type 512. *Antimicrob Agents Chemother*. 2016; 60(9):5612–5.
29. Yang YQ, Li YX, Song T, Yang YX, Jiang W, Zhang AY, et al. Colistin resistance gene *mcr-1* and its variant in *Escherichia coli* isolates from chickens in China. *Antimicrob Agents Chemother*. 2017; 61(5):e01204-16.
30. Zhao F, Feng Y, Lü X, McNally A, Zong Z. Remarkable Diversity of *Escherichia coli* Carrying *mcr-1* from Hospital Sewage with the Identification of Two New *mcr-1* Variants. *Front Microbiol*. 2017;8:2094.
31. Estabrook M, Wise M, Stone G, Kazmierczak K and Hackel M. Novel variant of transmissible colistin resistance determinant *mcr-1*. Unpublished

32. Lu X, Hu Y, Luo M, Zhou H, Wang X, Du Y, et al. *MCR-1.6*, a new *MCR* variant carried by an IncP plasmid in a colistin-resistant *Salmonella enterica* serovar Typhimurium isolate from a healthy individual. *Antimicrob Agents Chemother*. 2017; 61(5):e02632-16.
33. *Escherichia coli* strain WCHEC1604 phosphoethanolamine – lipid A transferase *MCR-1.7* (*mcr-1*) gene, *mcr-1.7* allele, complete cds. GenBank: KY488488.1.
34. *Escherichia coli* strain ST101 phosphoethanolamine transferase (*mcr-1.8*) gene, complete cds. GenBank: KY683842.1.
35. *Escherichia coli* strain HKSH_MCR_161114268_EC phosphoethanolamine lipid A transferase (*mcr1*) gene, *mcr1.9* allele, complete cds. GenBank: KY685071.1
36. Abuoun M, Stubberfield EJ, Duggett NA, Kirchner M, Dormer L, Nunez-Garcia J, et al. *mcr-1* and *mcr-2* (*mcr-6.1*) variant genes identified in *Moraxella* species isolated from pigs in Great Britain from 2014 to 2015. *J Antimicrob Chemother*. 2017;72(10):2745–9.
37. Partridge SR, Di Pilato V, Doi Y, Feldgarden M, Haft DH, Klimke W, et al. Proposal for assignment of allele numbers for mobile colistin resistance (*mcr*) genes. *J Antimicrob Chemother*. 2018;73(10):2625–30.
38. Current list of *mcr* gene families:
https://www.eurl-ar.eu/CustomData/Files/Folders/25-resourcer/554_colistin-all-genes-19-06-20.pdf
39. Xavier BB, Lammens C, Ruhel R, Kumar-Singh S, Butaye P, Goossens H, et al. Identification of a novel plasmid-mediated colistin-resistance gene, *mcr-2*, in *Escherichia coli*, Belgium, June 2016. *EuroSurveill*. 2016;21(27).
40. Garcia-Graells C, De Keersmaecker SCJ, Vanneste K, Pochet B, Vermeersch K, Roosens N, et al. Detection of plasmid-mediated colistin resistance, *mcr-1* and *mcr-2* genes, in *Salmonella* spp. isolated from food at retail in Belgium from 2012 to 2015. *Foodborne Pathog Dis*. 2018; 15(2):114–7.
41. Yin W, Li H, Shen Y, Liu Z, Wang S, Shen Z, et al. Novel plasmid-mediated colistin resistance gene *mcr-3* in *Escherichia coli*. *MBio*. 2017; 8(4):e01166-17.
42. Carattoli A, Villa L, Feudi C, Curcio L, Orsini S, Luppi A, et al. Novel plasmid-mediated colistin resistance *mcr-4* gene in *Salmonella* and *Escherichia coli*, Italy 2013, Spain and Belgium, 2015 to 2016. *EuroSurveill*. 2017;22(31).
43. Carretto E, Brovarone F, Nardini P, Russello G, Barbarini D, Pongolini S, et al. Detection of *mcr-4* positive *Salmonella enterica* serovar Typhimurium in clinical isolates of human origin, Italy, October to November 2016. *EuroSurveill*. 2018; 23(2).
44. Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, Malorny B. Identification of a novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar Paratyphi B. *J Antimicrob Chemother*. 2017; 72(12):3317–24.
45. Hammerl JA, Borowiak M, Schmoger S, Shamoun D, Grobbel M, Malorny B, et al. *mcr-5* and a novel *mcr-5.2* variant in *Escherichia coli* isolates from food and food-producing animals, Germany, 2010 to 2017. *J Antimicrob Chemother*. 2018; 73(5):1433–5.
46. Yang YQ, Li YX, Lei CW, Zhang AY, Wang HN. Novel plasmid-mediated colistin resistance gene *mcr-7.1* in *Klebsiella pneumoniae*. *J Antimicrob Chemother*. 2018;73(7):1791-1795.
47. *Klebsiella pneumoniae* strain SC20141012 plasmid pSC20141012, complete sequence. GenBank: MG267386.1
48. Wang X, Wang Y, Zhou Y, Li J, Yin W, Wang S, Zhang S, Shen J, Shen Z, Wang Y. Emergence of a novel mobile colistin resistance gene, *mcr-8*, in NDM-producing *Klebsiella pneumoniae*. *Emerg Microbes Infect*. 2018 Jul 4;7(1):122.
49. Carroll LM, Gaballa A, Guldemann C, Sullivan G, Henderson LO, Wiedmann M. Identification of Novel Mobilized Colistin Resistance Gene *mcr-9* in a Multidrug-Resistant, Colistin-Susceptible *Salmonella enterica* Serotype Typhimurium Isolate. *MBio*. 2019 May 7;10(3). pii: e00853-19.
50. Wang C, Feng Y, Liu L, Wei L, Kang M, Zong Z. Identification of novel mobile colistin resistance gene *mcr-10*. *Emerg Microbes Infect*. 2020 Mar 2;9(1):508-516.

51. Litrup E, Kiil K, Hammerum AM, Roer L, Nielsen EM, Torpdahl M. Plasmid-borne colistin resistance gene *mcr-3* in *Salmonella* isolates from human infections, Denmark, 2009–17. *Euro Surveill.* 2017;22(31).
52. Hernández M, Iglesias MR, Rodríguez-Lázaro D, Gallardo A, Quijada N, Miguela-Villoldo P, et al. Co-occurrence of colistin-resistance genes *mcr-1* and *mcr-3* among multidrug-resistant *Escherichia coli* isolated from cattle, Spain, September 2015. *Euro Surveill.* 2017;22(31).
53. Rebelo AR, Bortolaia V, Kjeldgaard JS, Pedersen SK, Leekitcharoenphon P, Hansen IM, et al. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro Surveill.* 2018; 23(6).
54. Liu L, Feng Y, Zhang X, McNally A, Zong Z. New Variant of *mcr-3* in an Extensively Drug-Resistant *Escherichia coli* Clinical Isolate Carrying *mcr-1* and *blaNDM-5*. *Antimicrob Agents Chemother.* 2017 Nov 22;61(12). pii: e01757-17.
55. Xiang R, Liu BH, Zhang AY, Lei CW, Ye XL, Yang YX, et al. Colocation of the polymyxin resistance gene *mcr-1* and a variant of *mcr-3* on a plasmid in an *Escherichia coli* isolate from a chicken farm. *Antimicrob Agents Chemother.* 2018; 62(6):e00501–18.
56. Karvanen M, Malmberg C, Lagerbäck P, Friberg LE, Cars O. Colistin is extensively lost during standard in vitro experimental conditions. *Antimicrob Agents Chemother.* 2017; 61:e00857–17.
57. EUCAST warning concerning antimicrobial susceptibility testing products or procedures. European Committee on Antimicrobial Susceptibility Testing; 2018 (http://www.eucast.org/ast_of_bacteria/warnings/).
58. Osei Sekyere J. Mcr colistin resistance gene: a systematic review of current diagnostics and detection methods. *Microbiology Open.* 2019;8(4):e00682
59. Simner PJ, Bergman Y, Trejo M, et al. Two-site evaluation of the colistin broth disk elution test to determine colistin in vitro activity against Gram-negative bacilli. *J Clin Microbiol.* 2019;57(2):e01163-18.
60. Routine internal quality control as recommended by EUCAST. Version 10.0. European Committee on Antimicrobial Susceptibility Testing; 2020.
61. Recommendations for MIC determination of colistin (polymyxin E). Joint CLSI-EUCAST Polymyxin Breakpoints Working Group. European Committee on Antimicrobial Susceptibility Testing; 2016 (http://www.eucast.org/guidance_documents/).
62. Breakpoint tables for interpretation of MICs and zone diameters. Version 10.0. European Committee on Antimicrobial Susceptibility Testing; 2020.
63. Satlin MJ, Lewis JS, Weinstein MP, et al. Clinical and Laboratory Standards Institute (CLSI) and European Committee on Antimicrobial Susceptibility Testing (EUCAST) position statements on polymyxin B and colistin clinical breakpoints. *Clin Infect Dis.* 2020;ciaa121.
64. Van Duin D, Lok JJ, Earley M, Cober E, Richter SS, Perez F, et al. Colistin versus ceftazidime-avibactam in the treatment of infections due to carbapenem-resistant *Enterobacteriaceae*. Antibacterial Resistance Leadership Group. *Clin Infect Dis.* 2018; 66(2):163–71.
65. Norgaard SM, Jensen CS, Aalestrup J, Vandenbroucke-Grauls CMJE, de Boer MGJ, Pedersen AB. Choice of therapeutic interventions and outcomes for the treatment of infections caused by multidrug-resistant gram-negative pathogens: a systematic review. *Antimicrob Resist Infect Control.* 2019 Nov 4;8:170.
66. Nation RL, Garonzik SM, Thamlikitkul V, Giamarellos-Bourboulis EJ, Forrest A, Paterson DL, et al. Dosing guidance for intravenous colistin in critically-ill patients. *Clin Infect Dis.* 2017; 64(5):565–71.
67. Abdul Momin MHF, Bean DC, Hendriksen RS, Haenni M, Phee LM, Wareham DW. CHROMagar COL-APSE: a selective bacterial culture medium for the isolation and differentiation of colistin-resistant Gram-negative pathogens. *J Med Microbiol.* 2017; 66:1554–61.
68. Nordmann P, Jayol A, Poirel L. A universal culture medium for screening polymyxin-resistant Gram-negative Isolates. *J Clin Microbiol.* 2016;54(5):1395–9

69. CHROMID® Colistin R Agar (COLR)
<https://www.biomerieux-diagnostics.com/chromid-colistin-r>
70. Nordmann P, Jayol A, Poirel L. Rapid detection of polymyxin resistance in *Enterobacteriaceae*. *Emerg Infect Dis*. 2016; 22(6):1038–43.
71. Boyen F, Vangroenweghe F, Butaye P, De Graef E, Castryck F, Heylen P, et al. Disk prediffusion is a reliable method for testing colistin susceptibility in porcine *E. coli* strains. *Vet Microbiol*. 2010; 144(3–4):359–62.
72. Matuschek E, Åhman J, Webster C, Kahlmeter G. Antimicrobial susceptibility testing of colistin – evaluation of seven commercial MIC products against standard broth microdilution for *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Acinetobacter* spp. *Clin Microbiol Infect*. 2018 Aug; 24(8):865–870.
73. Wayne, PA. CLSI. Performance Standards for Antimicrobial Susceptibility Testing. 30th ed. CLSI supplement M100. Clinical and Laboratory Standards Institute; M100 30th edition 2020
74. Humphries RM, Green DA, Schuetz AN, Bergman Y, Lewis S, Yee R, Stump S, Lopez M, Macesis N, Uhlemann AC, Kohner P, Cole N, Simner PJ. Multicenter Evaluation of Colistin Broth Disk Elution and Colistin Agar Test: a Report from the Clinical and Laboratory Standards Institute. *Journal of Clinical Microbiology* Oct 2019, 57 (11) e01269-19; DOI: 10.1128/JCM.01269-19
75. Pasteran F, Danze D, Menocal A, Cabrera C, Castillo I, Albornoz E, Lucero C, Rapoport M, Ceriana P, Corso A. Simple phenotypic tests to improve accuracy in screening chromosomal and plasmid-mediated colistin resistance in gram-negative bacilli. *bioRxiv* 2020.07.06.190850; doi: <https://doi.org/10.1101/2020.07.06.190850>
76. Global antimicrobial resistance surveillance system: manual for early implementation. Geneva: World Health Organization; 2015.
77. Terveer EM, Nijhuis RHT, Crobach MJT, Knetsch CW, Veldkamp KE, Gooskens J, Kuijper EJ, Claas ECJ. Prevalence of colistin resistance gene (*mcr-1*) containing *Enterobacteriaceae* in feces of patients attending a tertiary care hospital and detection of a *mcr-1* containing, colistin susceptible *E. coli*. *PLoS One*. 2017 Jun 2;12(6):e0178598.
78. Rebelo AR, Bortolaia V, Kjeldgaard JS, Pedersen SK, Leekitcharoenphon P, Hansen IM, Guerra B, Malorny B, Borowiak M, Hammerl JA, Battisti A, Franco A, Alba P, Perrin-Guyomard A, Granier SA, De Frutos Escobar C, Malhotra-Kumar S, Villa L, Carattoli A, Hendriksen RS. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro Surveill*. 2018 Feb;23(6).
79. Borowiak M, Baumann B, Fischer J, et al. Development of a Novel *mcr-6* to *mcr-9* Multiplex PCR and Assessment of *mcr-1* to *mcr-9* occurrence in colistin-resistant *Salmonella enterica* isolates from environment, feed, animals and food (2011–2018) in Germany. *Front Microbiol*. 2020;11:80.
80. Zavascki AP, Girardello R, Magagnin CM, Antchevis LC, Maciel RA, Palmeiro JK, et al. Emergence of polymyxin B resistance in a polymyxin B-susceptible KPC-producing *Klebsiella pneumoniae* causing bloodstream infection in a neutropenic patient during polymyxin B therapy. *Diagn Microbiol Infect Dis*. 2018;90(2):134–8.
81. Zankari E, Hasman H, Kaas RS, Seyfarth AM, Agersø Y, Lund O, et al. Genotyping using whole-genome sequencing is a realistic alternative to surveillance based on phenotypic antimicrobial susceptibility testing. *J Antimicrob Chemother*. 2013; 68(4):771–7.
82. Hendriksen RS, Bortolaia V, Tate H, Tyson GH, Aarestrup FM, McDermott PF. Using Genomics to Track Global Antimicrobial Resistance. *Front Public Health*. 2019;7:242.
83. Overview of services: Lyngby: Center for Genomic Epidemiology; 2018 (<https://cge.cbs.dtu.dk/services/>).
84. ResFinder 4 (database updated on August 2019 to include all known *mcr*-genes and chromosomal point mutations <https://cge.cbs.dtu.dk/services/ResFinder-4.0/>)
85. García-Fernández S, García-Castillo M, Ruiz-Garbajosa P, Morosini MI, Bala Y, Zambardi G, et al. Performance of CHROMID® Colistin R agar, a new chromogenic medium for screening of colistin-resistant *Enterobacterales*. *Diagn Microbiol Infect Dis*. 2019; 93(1):1–4.
86. Girlich D, Naas T, Dortet L. Comparison of the Superpolymyxin™ and CHROMID® Colistin R screening media for the detection of colistin-resistant *Enterobacteriaceae* from spiked rectal swabs. *Antimicrob Agents Chemother*. 2018;63(1):e01618-18.



**World Health
Organization**

World Health Organization
20 avenue Appia
1211 Geneva 27 - Switzerland
<https://www.who.int/health-topics/antimicrobial-resistance>

